

its stages and had not taken the larva or pupa before. These latter are readily distinguished from those of other Coccinellids by their distinct yellowish appearance. I brought back several of the larvae, which in due course pupated and emerged satisfactorily.

Some Notes on Swiss Butterflies.

[Supplementary to the *Butterflies of Switzerland*, by Rev. Geo. Wheeler, M.A., F.E.S.]

By the late MR. A. J. FISON.

(Arranged and communicated by Miss L. M. FISON.)

"The works of the Lord are great, sought out of all them that have pleasure therein."—*Psaln*, cxi. 2.

(Concluded from vol. xxxiv., page 217.)

Epinephele lycaon, Rott.—Fionnay, vii.; Nessel, 30.vii.90; Derborence, 25.viii.90.

Aphantopus hyperantus, L.—Rhone Bridge of St. Triphon, 21.viii.03 (lots); Charpigny, 3.viii.88; Gryonne, 22.vii.22; Champéry, 31.vii.22; Mt. Carré, vi. and viii.; Dent du Midi.

A. hyperantus ab. *arete*, Milliére.—Above Plan Cerisier; Martigny Combe.

A. hyperantus ab. *caeca*, Fuchs.—Corbeyrier; Éclépens.

Coenonympha iphix, Schiff.—Bex; Montet, 15.vi.20; Gryonne Meadows, 1.vii.22; Charpigny, 3.vii.22; St. Triphon Marsh, 3.vii.22; Dent du Midi, 4.vii.22; Mont Carré, 21.vi.

C. arcania, L., var. *insubrica*, Rätzer.—Le Prese, 13.vii.01; Faido, 13.vii.03 (1).

C. arcania var. *darwiniana*, Stdgr.—Bel Alp, 14.vi.90; Caux, vi.

C. satyrion, Esp.—Dent du Midi; Barmaz; Bonaveau, 8.viii.22 (L.M.F.), very small spots, some none.

C. satyrion, Esp., var. *unicolor*, Wh.—Mortersatsch Glacier, 19.vii.01; Émosson; Col de Forelaz, 7.viii.09; Téméley, 2.viii.10. No black spots on band.

C. satyrion ab. *caeca*, Wh.—(1) Rigi Staffelhöhe, 21.vi.04.

C. pamphilus var. *lyllus*, Esp.—Mt. Bré; Cassarate; Mendrisio, 15.viii.02.

C. tiphon, Rott.—Altmatt, 17.vii.03; Arth-Goldau, 20.vi.04; Einsiedeln, 6.vii.04; Aigle beyond golf links by Canal, 16.vi.05, S.E. corner, the best part. The forms found there are very interesting, some being dark and some pale. Leeuk, 19.vii.07.

Oeneis aello, Hb.—Saasgrund, 29.vi.94; under mountains from Vernayaz to Martigny, no doubt blown down. Laquinthal, 18.vii.08; under Dent de Moreles, 25.vi.90; Faulhorn, 1.viii.14 (L.M.F.).

Erebia epiphron, Knoch., var. *cassiope*, Fabr.—Under Bel Alp, 14.vii.90.

E. epiphron ab. *nelamus*, Bdv.—Furka; Trübsee; Zermatt; Simplon; St. Bernard; Glacier du Trient.

E. epiphron var. *valesiana*, Mey.D.—Pierre-à-Voir; Mägenwald.

E. melampus, Fssl., ab. *sudetica*, Stgr.—Randa; (1) Prätigau.

E. flavofasciata, Heyne.—Schafberg, S. of 1st Restaurant, 1-15.vii.04; Tscherva glacier, 30 mins. below and behind Restaurant, 1 ♂, 10.vii.01; none there in 1904; first found about 1850. Now on S. side of Albrun Pass.

E. eriphyle, Frr.—St. Lucia, 30.vii.97.

E. pharte, Hb., ab. *phartina*, Stgr.—Anthémoz, 29.vii.02.

E. mnestra, Hb.—Boden Glacier, 18.vii.95.

E. alecto, Hb., var. *glacialis*, Esp.—Fully; Portailles; Vernayaz, 18.vii.91; Mattmark See, 29.vi.94; Faulhorn, 1.viii.14 (L.M.F.); Anzeindaz, 14.vii.91.

E. manto, Esp.—Dischmathal (near *eriphyle* locality), 11.vii.04; Fénistral Bridge, 4.viii.10; Chamossaire; Corbeyrier; Rochers de Naye; varies greatly in size.

E. manto, Esp., ab. *caecilia*, Hb.—Trient glacier.

E. ceto, Hb.—Belalp, 14.vii.90; Teméley, 8.viii.1900.

E. medusa, Fabr.—With extra eye at apex of forewing (on one side only) like ab. *eriades*, Wh. Found 1 in 50 soat Charpigny; and on the West side of Gryonne between road and railway in May, 1906, I got about 15 so. ♀s seem to have this eye at times. (18) Éclepens (marsh), 20.v.90; Belalp, 14.vii.90; Monte Bré, 28.v.03; Caux, 18.vi.03; Faido, 11.vi.04; (3) Charpigny, 8.vi.07.

E. oeme, Hb.—Maderanerthal, 24.vi.96; Faulhorn, 1.viii.14; Dent du Midi, 29.vi.14, with var. *valesiaca*, Elwes; Lutschine Valley, 3.viii.14; Champéry, 31.vii.22; Barmaz, 4.viii.22 (L.M.F.).

E. erias, Godt.—Losone (1), 23.v.03; Asegna, 4.v.96; above Inden, 4.vi.94; Dent de Morcles (2/3 up), 15.v.94.

E. nerine, Frr., var. *stelviana*, Curó.—Under Piz Languard (near 1st and 2nd snow), vii.04; Täschthal, 9.vii.95.

E. euryale, Esp.—Émaney, 13.viii.09.

E. liqea, L.—Dent du Midi, 29.vi.14—21.vii.22; Gorge de Chauderon; E. of Niouc, vii.; Frenières, 3.vii.13; Grindelwald, viii.14; Champéry, 30.vii.22 (L.M.F.).

E. aethiops, Esp.—Charpigny, viii.

E. aethiops ab. *leucotaenia*, Stgr.—Alpe de Bodine.

Erebia pronöe, Esp., var. *pitho*, Hb.—Salanfe Road above Salvan, 3.ix.10.

E. goante, Esp.—Émosson (Finhaut), 12.viii.09; sous Kippel, 10.viii.90; Faulhorn, 1.viii.14 (L.M.F.).

E. gorge, Esp.—Rothhorn nr. Brienz, 16.vii.92; Mattmark See, 29.vi.94; Anzeindaz, 14.vii.91; Gornergrat, 5.vi.97.

E. tyndarus, Esp.—Faulhorn, 1.viii.14; Barmaz, 4.viii.22 (L.M.F.).

E. tyndarus var. *dromus*, H.S.—Nr. Anthémoz, 4.viii.03; Barmaz, 1.viii.03.

E. lappona, Esp.—Arpilles, nr. Martigny (2082 m.) (4), 9.vi.90; Faulhorn, 1.viii.14 (L.M.F.).

Melanargia galathea, L.—Charpigny, vi.; Les Posses; nr. Gryonne at Bex. Some with blue spots at edge of W. Sierre, 8.vii.02; Bué, 3.vii.01; Bex, 8.vi.03.

M. galathea var. *galaxaera*, Esp.—Lavorgo, 9.vii.03; Mt. Bré, 28.v.03 and 18.viii.02; (24) Lugano, 2.vi.96.

NOTES ON COLLECTING, etc.

WARWICKSHIRE RECORDS OF ICHNEUMONIDAE (ICHNEUMONINAE).—Considering the meagre reports of *Ichneumonidae* from the Midlands, perhaps the following captures in the Coventry district will be of